


THE UNCROSSING PARTIAL ORDER ON MATCHINGS IS EULERIAN

THOMAS LAM


ABSTRACT. We prove that the partial order P_n on the set of matchings of $2n$ points on a circle, given by resolving crossings, is an Eulerian poset.

1. THE THEOREM

Let P_n denote the set of matchings on $2n$ points, labeled $1, 2, \dots, 2n$ in order on a circle. Each $\tau \in P_n$ can be represented by (usually many) medial graphs, or strand diagrams. We say that a medial graph is *lensless* if any two strands intersect at most once. The following lensless medial graph represents $\{(1, 7), (2, 9), (3, 8), (4, 10), (5, 6)\} \in P_5$.



For $\tau \in P_n$ we let $c(\tau)$ denote the number of crossings in a lensless medial graph representing τ . The set P_n can be equipped with a partial order obtained by resolving crossings:




We declare $\tau' \leq \tau$ if there is a lensless medial graph G representing τ such that resolving a crossing in G gives a lensless medial graph G' representing τ' . The partial order P_n is the transitive closure of these cover relations.

The poset P_n is graded with rank function given by $c(\tau)$, and we refer the reader to [ALT, Ken, Lam] for alternative descriptions of P_n . The poset P_n has a unique maximum element, and Catalan number C_n of minimum elements. Let $\hat{P}_n$ denote P_n with a minimum $\hat{0}$ adjoined, where we declare that $c(\hat{0}) = -1$. Recall that a graded poset P with a unique minimum and a unique maximum, is *Eulerian* if, for every interval $[x, y] \subset P$ where $x < y$, the number of elements of odd rank in $[x, y]$ is equal to the number of elements of even rank in $[x, y]$. The following result was expected by many experts.

T.L. was supported by NSF grant DMS-1160726.

Theorem 1. $\hat{P}_n$ is an Eulerian poset.

Here is a picture of $\hat{P}_3$:



The same result for the Bruhat order of a Weyl group was established by Verma [Ver], and our approach is similar. Indeed, we rely on an embedding [Lam] of P_n into the (dual of the) Bruhat order of an affine symmetric group. Theorem 1 further amplifies the analogy between Bruhat order and the uncrossing partial order P_n .

Like Bruhat order, the poset P_n has a topological interpretation. In [Lam] a compactification E_n of the space of circular planar electrical networks with n boundary points was constructed. We have a stratification $E_n = \bigsqcup_{\tau \in P_n} E_\tau$ by *electroid cells*, and $\overline{E_\tau} = \bigsqcup_{\tau' \leq \tau} E_{\tau'}$. The same partial order occurs in the study of the positive orthogonal Grassmannian and scattering amplitudes for ABJM [HW, HWX]. The partial order P_n is also discussed in this context by Kim and Lee [KL] who observed a special case of Theorem 1. It seems to be expected by experts that these spaces are homeomorphic to balls.

We remark that in [ALT], it is shown that P_n is thin (intervals of length two are diamonds). The following conjecture is probably widely expected (see the related [ALT, Conjecture 3.3]):

Conjecture 1. $\hat{P}_n$ is lexicographically shellable.

An *EL*-shelling of $\hat{P}_3$ is given as follows. Denote the five non-crossing matchings by $\alpha_1, \alpha_2, \beta_1, \beta_2, \beta_3$, in order from left to right, as shown in the diagram above. We label each edge by one of the symbols $\{\alpha_1, \alpha_2, \beta_1, \beta_2, \beta_3\}$. Label the covers $\hat{0} \lessdot \tau$ by τ . Label the cover $\eta \lessdot \tau$ where $c(\tau) = 1$, by η' where $\eta' \neq \eta$ is uniquely defined by $\hat{0} \lessdot \eta' \lessdot \tau$. Label the cover $\eta \lessdot \tau$ where $c(\tau) = 2$, by β_i , where β_i is uniquely determined by the conditions $\beta_i < \tau$ and $\beta_i \not\leq \eta$. Label the cover $\eta \lessdot \hat{1} = \tau_{\text{top}}$ by β_i where β_i is uniquely determined by the condition $\beta_i \not\leq \eta$.




FIGURE 1. The Yang-Baxter move



FIGURE 2. Lens removal



FIGURE 3. Loop removal

Finally, order the symbols by $\alpha_1 < \beta_1 < \beta_2 < \beta_3 < \alpha_2$.

Acknowledgments. It is a pleasure to thank Yu-tin Huang, Rick Kenyon, and Pavlo Pylyavskyy for conversations motivating this work.

2. MEDIAL GRAPHS WITH LENSES

The notion of a medial graph representing a matching requires some care in the case that lenses are present.

Suppose G is any medial graph. A *lens* in G is a pair of strands that intersect twice or more. By repeatedly applying the moves in Figures 1 - 3, we can reduce G to a lensless medial graph G' . If G' represents a matching $\tau \in P_n$, we shall say that G represents $\tau \in P_n$ as well. The following result follows from the interpretation of medial graphs in terms of electrical networks [ALT, Ken, Lam].

Lemma 1. *Suppose G is a medial graph with lenses and G' , G'' are two lensless graphs obtained from G by the moves Figures 1 - 3. Then G' and G'' represent the same matching $\tau \in P_n$.*

Lemma 2. *Suppose G is a medial graph, possibly with lenses, that represents τ . Suppose G' is obtained from G by resolving any number of crossings in any way, and removing all closed interior loops that result. If G' represents τ' , then $\tau \geq \tau'$.*

Proof. In the language of [Lam], G is the medial graph of some electrical network Γ , such that $L(\Gamma) \in E_\tau$. Then G' is the medial graph of Γ' , where Γ' is obtained from Γ by either removing some edges, or contracting some edges. (Interior loops correspond to isolated interior components of Γ' .) Thus we must have $L(\Gamma') \in \overline{E_\tau}$, and the result follows from [Lam, Theorem 5.7], which states that $\overline{E_\tau} = \bigsqcup_{\eta \leq \tau} E_\eta$. $\square$

3. THE PROOF

For a subset $S \subset \hat{P}_n$, write $\chi(S) = \sum_{\tau \in S} (-1)^{c(\tau)}$. We need to show that $\chi([\tau, \eta]) = 0$ whenever $\tau < \eta$.

Let $\tilde{S}_{2n}$ denote the poset of affine permutations with period $2n$. In [Lam] we showed that there is an injection $\iota : P_n \hookrightarrow \tilde{S}_{2n}$ expressing P_n as dual to an induced subposet of $\tilde{S}_{2n}$. Write $g_\tau := \iota(\tau)$. We refer the reader to [Lam] for full details.

For $f \in \tilde{S}_{2n}$, we let

$$D_L(f) := \{i \in \mathbb{Z}/2n\mathbb{Z} \mid s_i f < f\} \quad \text{and} \quad D_R(f) := \{i \in \mathbb{Z}/2n\mathbb{Z} \mid f s_i < f\}$$

be the left and right descent sets of f . The following result is standard, see [BB, Proposition 2.2.7].

Lemma 3. *Suppose $f \leq g$ in $\tilde{S}_{2n}$. If $i \in D_L(g) \setminus D_L(f)$ then $f \leq s_i g$ and $s_i f \leq g$. If $i \in D_R(g) \setminus D_R(f)$ then $f \leq g s_i$ and $f s_i \leq g$.*

For $\tau \in P_n$ and $i \in \mathbb{Z}/2n\mathbb{Z}$, we let

$$i \in \begin{cases} A(\tau) & \text{if the strands labeled } i \text{ and } i+1 \text{ do not cross,} \\ B(\tau) & \text{if the strands labeled } i \text{ and } i+1 \text{ cross,} \\ C(\tau) & \text{if } i \text{ is matched with } i+1, \end{cases}$$

so that we have a disjoint union $\mathbb{Z}/2n\mathbb{Z} = A(\tau) \cup B(\tau) \cup C(\tau)$. Define $s_i \cdot \tau$ by

$$g_{s_i \cdot \tau} = s_i g_\tau s_i.$$

For example, if $i \in A(\tau)$ then $s_i \cdot \tau$ is obtained from τ by adding a crossing between strands i and $i+1$, close to the boundary at i and $i+1$. Note that we have

$$i \in \begin{cases} A(\tau) & \text{if } s_i g_\tau s_i < g_\tau \text{ or, equivalently, } s_i \cdot \tau \succ \tau, \\ B(\tau) & \text{if } s_i g_\tau s_i > g_\tau \text{ or, equivalently, } s_i \cdot \tau \prec \tau, \\ C(\tau) & \text{if } s_i g_\tau s_i = g_\tau \text{ or, equivalently, } s_i \cdot \tau = \tau. \end{cases}$$

Lemma 4. *Suppose $\tau \leq \eta$ in P_n . If $i \in A(\tau) \cap B(\eta)$ then $\tau \leq s_i \cdot \eta$ and $s_i \cdot \tau \leq \eta$.*

Proof. Since $i \in A(\tau)$ we have $i \in D_L(g_\tau) \cap D_R(g_\tau)$. Since $i \in B(\eta)$ we have $i \notin D_L(g_\eta) \cup D_R(g_\eta)$. By Lemma 3, we have $s_i g_\eta \leq g_\tau$. We also have, $i \notin D_R(s_i g_\eta)$ since $g_{s_i \cdot \eta} = s_i g_\eta s_i > s_i g_\eta$. So by Lemma 3 again, we have $s_i g_\eta s_i \leq g_\tau$, or equivalently, $g_{s_i \cdot \eta} \leq g_\tau$, or equivalently, $\tau \leq s_i \cdot \eta$. The proof of $s_i \cdot \tau \leq \eta$ is similar. $\square$

The following result follows easily from the “uncrossing” definition.

Lemma 5. *Suppose $\tau \leq \eta$ and $i \in A(\tau)$. Then $i \notin C(\eta)$.*

Proof of Theorem 1. We shall first prove the theorem for intervals $\tau < \eta$ of P_n ; that is, intervals not involving $\hat{0}$. We proceed by descending induction on $c(\tau) + c(\eta)$. The base case where η is the maximal element of P_n and $c(\tau) = \binom{n}{2} - 1$ is clear. Also if $c(\eta) - c(\tau) = 1$ the result is clear, so we may assume that $c(\eta) - c(\tau) \geq 2$.

Since τ is not the maximal element, $D_L(g_\tau)$ and $D_R(g_\tau)$ are non-empty, so $A(\tau)$ is non-empty (see also the proof of [Lam, Lemma 4.14]). Let $i \in A(\tau)$.

Case 1: $i \in A(\eta)$. Then

$$(1) \quad [\tau, \eta] = [\tau, s_i \cdot \eta] \setminus \{\sigma \mid \tau \leq \sigma \leq s_i \cdot \eta, \sigma \not\leq \eta\}.$$

We claim that

$$(2) \quad \{\sigma \mid \tau \leq \sigma \leq s_i \cdot \eta, \sigma \not\leq \eta\} = \{\sigma \mid s_i \cdot \tau \leq \sigma \leq s_i \cdot \eta, \sigma \not\leq \eta\}$$

Suppose $\tau \leq \sigma \leq s_i \cdot \eta$ and $\sigma \not\leq \eta$. If $i \in A(\sigma)$, then since $i \in B(s_i \cdot \eta)$, applying Lemma 4 to $\sigma < s_i \cdot \eta$ we would obtain $\sigma \leq \eta$, contradicting our assumption. If $i \in C(\sigma)$, then Lemma 5 would be violated. Thus $i \in B(\sigma)$, or $s_i \cdot \sigma < \sigma$. Now apply Lemma 4 to $\tau \leq \sigma$ to obtain $s_i \cdot \tau \leq \sigma$. This proves (2).

But we have

$$\{\sigma \mid s_i \cdot \tau \leq \sigma \leq s_i \cdot \eta, \sigma \not\leq \eta\} = [s_i \cdot \tau, s_i \cdot \eta] \setminus [s_i \cdot \tau, \eta]$$

so by induction $\chi(\{\sigma \mid s_i \cdot \tau \leq \sigma \leq s_i \cdot \eta, \sigma \not\leq \eta\}) = \chi([s_i \cdot \tau, s_i \cdot \eta]) - \chi([s_i \cdot \tau, \eta]) = 0$. By induction again, $\chi([\tau, s_i \cdot \eta]) = 0$ so using (1) we obtain $\chi([\tau, \eta]) = 0$.

Case 2: $i \in B(\eta)$. We apply Lemma 4 to see that $\sigma \in [\tau, \eta]$ implies $s_i \cdot \sigma \in [\tau, \eta]$. By Lemma 5, it follows that $\sigma \mapsto s_i \cdot \sigma$ is an involution which swaps elements of odd rank with elements of even rank. Thus $\chi([\tau, \eta]) = 0$.

Case 3: $i \in C(\eta)$. This is impossible by Lemma 5.

Thus we have shown that $\chi([\tau, \eta]) = 0$ for intervals $\tau < \eta$ not involving $\hat{0}$.

Now suppose $\tau = \hat{0}$ and $\hat{0} < \eta$. We may suppose that $c(\eta) \geq 1$. Then $B(\eta) \neq \emptyset$. Let $i \in B(\eta)$. By Lemma 4, the map $\sigma \mapsto s_i \cdot \sigma$ establishes an involution on the set $\{\sigma \in (\hat{0}, \eta] \mid i \notin C(\eta)\}$, and this involution swaps the parity of $c(\sigma)$. Let

$$S = \{\sigma \in (\hat{0}, \eta) \mid i \in C(\eta) \text{ or } \sigma = \hat{0}\}.$$

It thus suffices to show that $\chi(S) = 0$. We claim that S has a unique maximal element κ . This would complete the proof since then $S = [\hat{0}, \kappa]$ and we may proceed by induction.

We now construct $\kappa \in P_n$. Let G be a medial graph representing η . The strands p_i and p_{i+1} beginning at i and $i+1$ cross each other since $i \in B(\eta)$. We shall assume p_i and p_{i+1} cross each other at q before intersecting any other strands. Let G' be obtained from G by uncrossing q so that i is matched directly with $i+1$. Let $\kappa \in P_n$ be represented by G' .

Suppose $\sigma \in S$. We need to show that $\sigma \leq \kappa$. A (lensless) medial graph G'' representing σ can be obtained from G by uncrossing some subset $\mathcal{C}$ of the crossings of G (see the comment after [Lam, Lemma 4.11]). Since $i \in C(\sigma)$ we must have $q \in \mathcal{C}$. If q is resolved in G'' in the same way as in G' then we are done: G'' can be obtained from G' by a number of uncrossings, and Lemma 2 gives $\sigma \leq \kappa$. Now suppose q is resolved in G'' in the direction different to the one in G' . We then observe that the medial graph G''' obtained from G'' by resolving the uncrossing q in the other direction also represents the same matching σ (a closed interior loop will appear in G''' , which can be removed). This completes the proof that $S = [0, \kappa]$, and thus the theorem. $\square$

REFERENCES

- [ALT] J. ALMAN, C. LIAN, AND B. TRAN: Circular Planar Electrical Networks I: The Electrical Poset EP_n , preprint, 2013; [arXiv:1309.2697](#).
- [BB] A. BJÖRNER AND F. BRENTI: Combinatorics of Coxeter groups. Graduate Texts in Mathematics, 231. Springer, New York, 2005. xiv+363 pp.
- [HW] Y.-T. HUANG AND C. WEN: ABJM amplitudes and the positive orthogonal grassmannian, preprint, 2013; [arXiv:1309.3252](#).
- [HWX] Y.-T. HUANG, C. WEN, AND D. XIE: The Positive orthogonal Grassmannian and loop amplitudes of ABJM, preprint, 2014; [arXiv:1402.1479](#).
- [Ken] R. KENYON: The Laplacian on planar graphs and graphs on surfaces, preprint; [arXiv:1203.1256](#).
- [KL] J. KIM AND S. LEE: Positroid Stratification of Orthogonal Grassmannian and ABJM Amplitudes, preprint, 2014; [arXiv:1402.1119](#).
- [Lam] T. LAM: Electroid varieties and a compactification of the space of electrical networks, preprint, 2014; [arXiv:1402.6261](#).
- [Ver] D.N. VERMA: Möbius inversion for the Bruhat ordering on a Weyl group, Ann. Sci. École Norm. Sup. 4 (1971), 393–398.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MICHIGAN, 2074 EAST HALL, 530 CHURCH STREET, ANN ARBOR, MI 48109-1043, USA

E-mail address: tfylam@umich.edu